

ARTICLE

HLA Class II Polymorphism: Protective or Risk Factors to Breast Cancer in Tunisia?

Amal BACCAR HARRATH,¹ Besma YACOUBI LOUESLATI,¹ Wafa TROUDI,¹ Slama HMIDA,² Syrine SEDKAOUI,¹ Amel DRIDI,² Afef JRIDI,² Farhat BEN AYED,³ Khaled BEN RHOMDHANE,³ Amel BEN AMMAR ELGAAIED¹

¹Laboratory of Molecular Genetics, Immunology and Biotechnology, Faculty of Sciences of Tunis, ElManar University, ²Tunisian Center of Blood Transfusion, ³Salah Azeiz Oncology Institute

HLA system plays a key role in the tumor cells' escape from immune surveillance. Herein is the first report on the correlation of the susceptibility to breast cancer with HLA class II markers in Tunisia. Molecular typing of HLA-DRB1 and -DQB1 loci was undertaken for 70 Tunisian female patients.

Comparison of allele and haplotype distribution between patients and 70 female control subjects reveals a negative association between HLA-DRB1*07-DQB1*02 and the incidence of breast cancer in the Tunisian population. (Pathology Oncology Research Vol 12, No 2, 79–81)

Key words: HLA, DRB1, DQB1, breast cancer

Introduction

The process of oncogenesis depends on the genetic instability of tumors and interactions with their immunological environment. Considering the importance of immune surveillance during tumorigenesis,^{13,17} some individuals who inherit specific alleles or haplotypes of the highly polymorphic HLA genes may be exposed or may resist to specific types of cancers.^{2,3,21,23,25} In fact, besides its crucial role in human clinical transplantation, the HLA system plays a key role in the mounting and recruiting of the cytotoxic T lymphocytes against tumor antigens that may contribute to tumor evasion from the destructive immune responses.

The major histocompatibility complex in humans, HLA, is a highly variable genetic system that is constituted by six main polymorphic loci, A, B, C, DRB1, DQB1 and DPB1, located on the short arm of chromosome 6.^{9,22} During the last decade, molecular typing techniques have been developed to identify HLA alleles at DNA level with a high-resolution power, and DNA mutations now allow the identification of a high number of alleles according to the

loci (121, 69, 429, 751, 219 and 429 alleles described in 2006 for the loci DPB1, DQB1, DRB1, B, C and A, respectively (www.ebi.ac.uk/imgt/hla/)).

Among the different HLA loci, DRB1 and DQB1 have been accurately tested at the molecular level since the 11th international histocompatibility workshop. The DRB1 and DQB1 loci are only separated by 85 kb on the chromosome, and almost no cases of recombination between these two loci have been reported in family studies. Therefore, HLA-DRB1 and DQB1 alleles generally exhibit a highly significant level of linkage disequilibrium.⁵

Until now, few reports have been carried out on the association of HLA antigens and the susceptibility to breast cancer.^{1,4,6,15,24} Some of these studies are limited to a small number and lack enough power to reach significance. Three reports have analyzed the polymorphism of HLA-class II genes in breast cancer as a genetic risk factor.^{8,10,14} The DRB1*11 allele was defined as protector in Caucasian population,¹⁰ while DRB1*12 was positively associated with breast cancer for Iranian women.¹⁴ Other studies examined single nucleotide polymorphisms (SNPs) in the HLA class III region.^{11,19,20}

In order to better understand the role played by the HLA system in the process of oncogenesis, it would be very interesting to search for an association between the HLA polymorphism genes and breast cancer risk in different populations. Such studies will allow us to detect the hid-

Received: Oct 3, 2005; *accepted:* March 18, 2006

Correspondence: Amal BACCAR HARRATH Laboratory of Molecular Genetics, Immunology and Biotechnology, Department of Biology, Faculty of Sciences of Tunis. 1060. Tunis, Tunisia. Tel: +216 96 35 26 24, fax: +216 71 821 702, E-mail: amal.baccar@laposte.net

den alleles that can be considered as risk factors in breast cancer development.

Our study is a detailed molecular analysis of HLA DQB1, DRB1 alleles and HLA DRB1-DQB1 haplotypes for Tunisian patients with breast cancer and ethnically matched controls.

Materials and Methods

Population samples

Blood samples were collected from 70 female patients with breast cancer admitted in Salah Azeiz Oncology Institute from April to August, 2001. The age of patients ranged between 27 and 67 years. Seventy healthy female blood donors from the Tunisian Center of Blood Transfusion were considered as a control group. All subjects were unrelated volunteers originated from Tunisia. Five milliliters of venous blood with EDTA, as anticoagulant, were collected from each subject.

DNA typing

Genomic DNA was extracted from peripheral blood according to the method of Maniatis.¹⁸ The second exon of both DRB1 and DQB1 loci and exon 3 of DQB1 were amplified by a polymerase chain reaction (PCR) using specific primers provided by Innolipa kits from Innogenetics (Ref: 80617 and 80337 for DRB1 and DQB1 loci, respectively). The HLA allelic polymorphism of these two loci was then analyzed by RDB hybridization^{7,16} using Innolipa kits from Innogenetics (Ref: 80635 and 80336 for DRB1 and DQB1 loci, respectively).

Statistical analysis

HLA-DRB1 and -DQB1 allele and haplotype frequencies were directly estimated by the following equation [$f = (n/N) \times 100$] where "n" is the number and "N" is the total number of alleles or haplotypes. Allelic and haplotype frequencies in cancer patients and controls were compared using Pearson's χ^2 test or Fisher's exact test (when the number of subjects in a cell is <5). Odds ratios are given with 95% confidence limits. The EPI INFO 6 package program was used for this statistics analysis.

Results

HLA-DRB1 and -DQB1 typing has been undertaken for 70 patients with breast cancer and 70 healthy matched donors. For the DRB1 loci, the results showed a total of 11 alleles in patients with breast cancer and 12 alleles in the matched controls (Table 1). For DQB1 locus only five alleles were found (Table 2). No significant correlation with incidence of breast cancer was observed for these two loci.

Table 1. HLA-DRB1* alleles in breast cancer patients and normal controls

DRB1 alleles	Patients N=70 (%)	Controls N=70 (%)	P value
DRB1*01	7 (5)	8 (5.71)	NS
DRB1*03	29 (20.71)	19 (13.57)	NS
DRB1*04	13 (9.28)	17 (12.14)	NS
DRB1*07	14 (10)	23 (16.42)	NS
DRB1*08	3 (2.14)	1 (0.71)	NS
DRB1*09	1 (0.71)	0	NS
DRB1*10	4 (2.85)	1 (0.71)	NS
DRB1*11	26 (18.57)	29 (20.71)	NS
DRB1*12	4 (2.85)	5 (3.57)	NS
DRB1*13	23 (16.42)	15 (10.71)	NS
DRB1*14	0	1 (0.71)	NS
DRB1*15	16 (11.42)	20 (14.28)	NS
DRB1*16	0	1 (0.71)	NS

Nominal value for comparison, P= 0.05; degree of freedom = 1; NS: not significant

Table 2. HLA-DQB1* alleles in breast cancer patients and normal controls

DQB1 alleles	Patients N= 70 (%)	Controls N=70 (%)	P value
DQB1*02	38 (27.14)	40 (28.57)	NS
DQB1*03	46 (31.96)	45 (32.14)	NS
DQB1*04	5 (3.57)	6 (4.28)	NS
DQB1*05	17 (12.14)	18 (12.85)	NS
DQB1*06	34 (24.28)	31 (22.14)	NS

Nominal value for comparison, P= 0.05; degree of freedom = 1; NS: not significant

The results of HLA-DRB1-DQB1 haplotypes are shown in Table 3. As indicated, a negative significant association exists between HLA-DRB1*07-DQB1*02 haplotype and the occurrence of the disease [9/140 (6.42%) haplotypes in patients, versus 22/140 (15.71%) in control group] (P= 0.013; OR= 0.37; 95% confidence interval: 0.15-0.88).

Conclusion

Our study is the first report on the correlation of the susceptibility to breast cancer with HLA class II markers in Tunisian patients. The results revealed a negative association between HLA-DRB1*07-DQB1*02 and the occurrence of breast cancer. This association could be linked to the role of HLA class II molecules in the anti-tumoral response, or explained by linkage disequilibrium between these markers and the susceptibility or protective genes.

In fact, recent research has focused on the functional characteristics and the biological significance of MHC class II gene products in relation to processing and presen-

Table 3. HLA-DRB1*-DQB1* haplotypes in breast cancer patients and normal controls

<i>DRB1-DQB1 haplotypes</i>	<i>Patients N=70 (%)</i>	<i>Controls N= 70 (%)</i>	<i>P value</i>
DRB1*01 -DQB1*05	6 (4.28)	8 (5.71)	NS
DRB1*01-DRB1*03	1 (0.71)	0	NS
DRB1*03-DQB1*02	27 (19.28)	18 (12.85)	NS
DRB1*03 -DQB1*04	2 (1.42)	0	NS
DRB1*03-DQB1*06	0	1 (0.71)	NS
DRB1*04 -DQB1*03	11 (7.85)	12 (8.57)	NS
DRB1*04 -DQB1*04	2 (1.42)	5 (3.57)	NS
DRB1*07 -DQB1*02	9 (6.42)	22 (15.71)	P = 0.013
DRB1*07 -DQB1*03	5 (3.57)	1 (0.71)	NS
DRB1*08 -DQB1*03	2 (1.42)	0	NS
DRB1*08 -DQB1*04	1 (0.71)	1 (0.71)	NS
DRB1*09 -DQB1*02	1 (0.71)	0	NS
DRB1*10 -DQB1*05	4 (2.85)	1 (0.71)	NS
DRB1*11 -DQB1*03	22 (15.71)	22 (15.71)	NS
DRB1*11 -DQB1*05	4 (2.85)	7 (5)	NS
DRB1*12 -DQB1*03	2 (1.42)	5 (3.57)	NS
DRB1*12 -DQB1*05	2 (1.42)	0	NS
DRB1*13 -DQB1*02	1 (0.71)	0	NS
DRB1*13 -DQB1*03	3 (2.14)	5 (3.57)	NS
DRB1*13 -DQB1*06	19 (13.57)	10 (7.14)	NS
DRB1*14 -DQB1*05	0	1 (0.71)	NS
DRB1*15 -DQB1*05	1 (0.71)	0	NS
DRB1*15 -DQB1*06	15 (10.71)	20 (14.28)	NS
DRB1*16 -DQB1*05	0	1 (0.71)	NS

Nominal value for comparison, P= 0.05; degree of freedom = 1; NS: not significant

tation of tumor antigens and their role in anticancer strategies against solid tumors. Indeed, Feinmesser and his coworkers have demonstrated that it is likely that the inherited individual specificities affect the quality and quantity of anti-tumor immune response.¹²

Our study reveals a new association between HLA markers and breast cancer, and proves that it would be worthwhile to investigate the association of these markers with breast cancer risk in different world populations.

It is important to note that although we have examined a reasonable number of women with breast cancer, clearly more extensive studies need to be conducted on a large sample of patients and controls, in order to confirm this association and to detect other important alleles or haplotypes that could be considered as risk factors in the occurrence or presentation of breast cancer.

Acknowledgements

We thank all blood donors and the women with breast cancer who participated in the present study. We are grateful to the staff of Salah Azeiz Oncology Institute for their help in the collection of blood samples. This work was supported by the Tunisian Ministry of Research, technology and development of competences.

References

1. Abrahamova J, Majsky A: HLA system and some neoplastic diseases. *Acta Univ Carol Med Monogr* 123: 1-80, 1988
2. Apple RJ, Erlich HA, Klitz W, et al: HLA DR-DQ associations with cervical carcinoma show papillomavirus-type specificity. *Nat Genet* 6:157-162, 1994
3. Bidwell JL, Soong TW, Raymond PA, et al: HLA genotyping of colorectal carcinoma in the Chinese population. *Hum Immunol* 34:19-23, 1992
4. Biswal BM, Kumar R, Julka PK, et al: Human leucocytic antigens (HLA) in breast cancer. *Int J Med Sci* 52:177-183, 1998
5. Bodmer WF: The HLA system: structure and function. *J Clin Pathol* 40:948-958, 1987
6. Bouillence C, Deneufbourg JM: Positive correlation between breast cancer incidence and HLA antigens. *Oncology* 36:156-159, 1979
7. Buysse I, Dacorte R, Baens M, et al: Rapid DNA typing of class II antigens using the polymerase chain reaction and reverse dot blot hybridization. *Tissue Antigens* 41:1-14, 1993
8. Casoli C, Zanelli P, Adorni A, et al: Serological and molecular study on the HLA phenotype of female breast cancer patients. *Eur J Cancer* 30: 1207-1208, 1994
9. Campbell RD, Trowsdale J: Map of the human MHC. *Immunol Today* 14 349, 1993
10. Chaudhuri S, Cariappa A, Tang M, et al: Genetic susceptibility to breast cancer, HLA DQB*03032 and HLA DRB1*11 may represent protective alleles. *Proc Natl Acad Sci USA* 97: 11451-11454, 2000
11. De Jong MM, Nolte IM, De Vries EGE, et al: The HLA class III sub-region is responsible for an increased breast cancer risk. *Hum Mol Genet* 12:2311-2319, 2003
12. Feinmesser M, Sulkes A, Morgenstern S, et al: HLA-DR and beta 2 microglobulin expression in medullary and atypical medullary carcinoma of the breast: histopathologically similar but biologically distinct entities. *J Clin Pathol* 53: 286-291, 2000
13. Ferguson TA, Green DR, Griffith TS: Cell death and immune privilege. *Int Rev Immunol* 21:153-172, 2002
14. Ghaderi A, Talei A, Ghareisi-Fard B, et al: HLA-DRB1 alleles and the susceptibility of Iranian patients with breast cancer. *Pathol Oncol Res* 7: 39-41, 2001
15. Hammond MG, Appadoo B, Brain P: HLA and cancer in South African Indians. *Tissue Antigens* 14:296-302, 1979
16. Hmida S, Gauthier A, Dridi A, et al: HLA class II polymorphism in Tunisians. *Tissue Antigens* 45: 63-68, 1995
17. Igney FH, Krammer PH: Death and anti-death: tumor resistance to apoptosis. *Nat Rev Cancer* 2:277-288, 2002
18. Maniatis T, Fritsch EF, Sambrook J: *Molecular Cloning. A Laboratory Manual*. Cold Spring Laboratory Press, Cold Spring Harbor NY, 1982
19. Mestiri S, Bouaouina N, Ahmed SB, et al: Genetic variation in the tumor necrosis factor-alpha promoter region and in the stress protein hsp 70-2: susceptibility and prognostic implications in breast carcinoma. *Cancer* 91: 672-678, 2001
20. Park KS, Mok JW, Ko HE, et al: Polymorphisms of tumor necrosis factors A and B in breast cancer. *Eur J Immunogenet* 29: 7-10, 2002
21. Pellegris G, Illeni MT, Vaglini M, et al: HLA antigens in malignant melanoma patients. *Tumori* 66: 51-58, 1980
22. Ragoussis J, Monaco A, Mockridge I, et al: Cloning of the HLA class II region in yeast artificial chromosomes. *Proc Natl Acad Sci USA* 88: 3753-3757, 1991
23. Sastre-Garau X, Loste MN, Vincent-Salomon A, et al: Decreased frequency of HLA-DRB1 13 alleles in Frenchwomen with HPV-positive carcinoma of the cervix. *Int J Cancer* 69: 159-164, 1996
24. Subira ML, Crisci CD, Zornoza G, et al: Breast cancer and histocompatibility antigens. *Allegrol Immunopathol (Madr)* 7: 411-416, 1979
25. Wu MS, Hsieh RP, Huang SP, et al: Association of HLA-DQB1*0301 and HLA-DQB1*0602 with different subtypes of gastric cancer in Taiwan. *Jpn J Cancer Res* 93:404-410, 2002